



Vaping and Public Health: Scientific Evidence, Therapeutic Controversies, and Regulatory Challenges in Adolescents

Carmen Fernández-Cáceres^{1,✉}, José Luis Sánchez-Gutiérrez^{2,✉}, Rodrigo Marín-Navarrete^{3,✉}

¹ Dirección General, Centros de Integración Juvenil, México.

² Departamento de Investigación Psicosocial y Documental, Centros de Integración Juvenil, México.

³ Dirección de Investigación y Enseñanza, Centros de Integración Juvenil, México.

Several types of electronic cigarettes or vapes are currently available on the market. The World Health Organization (WHO) classifies them into two categories: electronic nicotine delivery systems (ENDS) and electronic non-nicotine delivery systems (ENNDS) (WHO, 2016). Both have health implications, as they operate in a comparable way (WHO, 2020). Moreover, health alerts have been issued because some products labeled as ENNDS contain nicotine despite their packaging indicating otherwise (Guraka et al., 2024; Miller et al., 2021; Omaiye et al., 2017). These devices operate using a battery and a metal heating element that vaporizes a mixture of substances known as electronic liquids (e-liquids), producing aerosols that are inhaled and exhaled by the user, a practice known as vaping (Baldassarri, 2020).

The characteristics and composition of the aerosol produced by vapes vary depending on factors such as battery power, the chemical components present in e-liquids, the topography of consumption, and atomizer design (Williams & Talbot, 2019).

According to the specialized literature, four generations of electronic vaping devices exist. The first generation is shaped like a conventional cigarette, is disposable, and uses low-voltage batteries. The second generation features variable voltage batteries and pre-filled or refillable tanks enabling users to adjust the nicotine concentration. The third generation is characterized by larger capacity tanks, customizable coils, and the ability to replace atomizer components. Finally, the fourth generation includes more compact devices using interchangeable capsules or pods integrating both the e-liquid reservoir and the heating coil (Baldassarri, 2020; Starnini et al., 2025; Williams & Talbot, 2019).

The e-liquids used in these devices are typically made with a base of propylene glycol and vegetable glycerin (Starnini et al., 2025) acting as a carrier for a complex blend of chemical compounds and flavorings (Pennington & Hernandez, 2025), as well as nicotine, either in its free form or as nicotine salts (Harris, 2024).

The heating process creates carbonyl byproducts and other reactive compounds in the aerosol, the concentration of which depends on the e-liquid's formulation and the type of heating used (Harris, 2024). Contact between the liquid and the metal components of the atomizer and coil degradation contribute to the presence of metals and metalloids (Schmidt, 2024). Moreover, in recent years, a rise in nicotine concentration has been observed, increasing the addictive potential of these products (Diaz et al., 2025).

Despite the evidence presented above, the expert community proposes the use of electronic nicotine replacement therapy (ENRT) devices as a therapeutic tool to reduce withdrawal symptoms and promote smoking cessation. Positive results have been shown in pregnant women (Przulj et al., 2023) and in adults planning to quit smoking (Levet et al., 2023).

Moreover, ENRT devices have been observed to be more effective in achieving abstinence than ENNDS, nicotine patches, and other therapeutic treatments (Levet et al., 2023).

However, the use of these devices for therapeutic purposes requires strict policies and regulations to ensure controlled doses of nicotine, given that many devices contain higher concentrations than conventional cigarettes (Diaz et al., 2025). Good

Corresponding author:

Rodrigo Marín Navarrete. Centros de Integración Juvenil. Avenida San Jerónimo 372, Col. Jardines del Pedregal, Alcaldía Álvaro Obregón, C.P. 01900, Ciudad de México. Email: rodrigo.marin@cij.gob.mx



manufacturing practices with quality standards and monitoring of components and concentrations are also required (Guraka et al., 2024). Access must be restricted to people with nicotine dependence who use them for therapeutic purposes and under medical supervision (Peri & Eisenberg, 2024). Finally, greater investment in research is needed to evaluate the risks associated with their use in clinical contexts (Feeney et al., 2022).

At the same time, there is also a proposal to use vapes as a harm reduction measure for smokers who do not wish to quit nicotine. These smokers are willing to change their consumption method (from smoking to vaping) on the assumption that vaping presents a lower risk of developing medical conditions.

This proposal has prompted the scientific community to investigate its true impact on health (Ailbright et al., 2024).

In this regard, a systematic review with meta-analysis was recently published, indicating that the use of electronic devices minimally reduces the risk of developing conditions such as asthma, chronic obstructive pulmonary disease (COPD), and stomatological diseases. However, evidence indicates that the risks of developing cardiovascular diseases, strokes, and metabolic dysfunctions are like those of conventional cigarette consumption. Additionally, people with dual consumption (smoking both conventional and e-cigarettes) have a 20% to 41% increase in the risk of developing any type of related disease, compared to exclusive conventional smokers (Glantz et al., 2024).

According to Hatsukami & Carroll (2020), the use of vapes as a harm reduction measure is controversial. Although smokers who do not wish to quit can benefit from reducing their exposure to combustion toxins, the devices pose new risks to non-smokers and younger generations. In this regard, the WHO (2023, 2024a, 2025) has urged countries to strengthen their public policies for the regulation, prevention and protection of children and adolescents. It points out that the industry's marketing strategies represent a threat by using attractive flavors, bright colors, and animated characters. These campaigns reduce the perception of risk through promotions, gifts, sponsorships at youth events and accelerated launches of new products taking advantage of regulatory gaps.

In addition, they are displayed in convenience stores alongside sweets and snacks (WHO, 2024b; Gaiha et al., 2024).

Leading brands in the sector, such as JUUL, PAX, and PLOOM, have employed marketing strategies that present vaping as a modern lifestyle, targeting young audiences. This includes promotion

on social networks through memes and teenage slang (Czaplicki et al., 2020; Gaiha et al., 2024) and insertion in popular media such as television, film, and music videos (Wakefield et al., 2025). Vaping devices are also presented as technological innovations with attractive designs (Jackler et al., 2019).

Given that vaping is a relatively recent product, several authors have underlined the need to assess the magnitude of the phenomenon and its impacts on the youth population (Novak & Wang, 2024). In this regard, evidence indicates that one in five Latin American adolescents has used e-cigarettes (Izquierdo-Condoy et al., 2024) and that adolescents who vape are three times more likely to start smoking conventional cigarettes than those who do not (Golder et al., 2025). Other research reports reveal that chronic vaping in children and adolescents is associated with airway fibrosis or constrictive bronchiolitis (Hariri et al., 2022). It is known to increase the risk of developing nicotine addiction, as well as cognitive impairments (attention and memory).

It also disrupts the circadian rhythm, in addition to producing anxiety, depression, and emotional dysregulation symptoms (Abdulhay et al., 2025; Javed et al., 2022; Meehan et al., 2024). In general, evidence indicates that nicotine exposure in this age group increases the risk of developing mental disorders due to its impact on the cholinergic and dopaminergic systems during neurodevelopment (Reynolds et al., 2025).

In Mexico, although the sale of electronic vaping devices is prohibited, data from the National Health and Nutrition Survey (ENSANUT) show a worrying change. Between 2018 and 2023, combustible tobacco consumption among young people aged 15 to 19 years decreased from 9.8% to 5%, while the exclusive use of vapes increased from 1.3% to 6.6% (Barrera-Núñez et al., 2025). In parallel, records from Centros de Integración Juvenil (CIJ) show an increase of over 2300% in lifetime nicotine use in electronic devices. The number of cases rose from 72 (1.1%) in the first half of 2021 to 1,738 (15.9%) in the first half of 2025. This phenomenon is observed across almost the entire country, with a higher prevalence in men (68.2%) than women (31.8%). The average age of onset of use is 16.5 years ($SD = 6.4$), and in 7.2% of cases, vaping was the main reason for consultation (Centros de Integración Juvenil, 2025a). Additionally, CIJ conducted a multisite cross-sectional survey in approximately 773 middle and high schools nationwide, including a sample of nearly 115,000 students. Preliminary findings revealed that 28% reported using vapers (Centros de Integración Juvenil, 2025b).

Taken together, the available evidence underlines that vaping in adolescents cannot be considered a harmless alternative or an effective tool for harm reduction. The sustained increase in the prevalence of vaping anticipates an overdemand for specialized services for cessation and the care of associated medical consequences, which will increase the health-care burden and public cost in the coming years. In this context, it is essential to establish an inter-institutional, inter-sectoral agenda focused on developing innovative prevention, treatment, and public policy strategies. The aim of these strategies would be to counteract the technological and symbolic appeal of these devices to strengthen the protection of new generations against nicotine dependence.

REFERENCES

- Abdulhay, N., Wiener, R. C., Wen, S., Gibbs, B. B., & Bhandari, R. (2025). Mental health outcomes associated with electronic cigarette use, combustible tobacco use, and dual use among U.S. adolescents: Insights from the National Youth Tobacco Survey. *PLOS Mental Health*, 2(7), e0000370. <https://doi.org/10.1371/journal.pmen.0000370>
- Allbright, K., Villandre, J., Alexander, L. E. C., Zhang, M., Benam, K. H., Evankovich, J., Königshoff, M., & Chandra, D. (2024). The paradox of the safer cigarette: understanding the pulmonary effects of electronic cigarettes. *European Respiratory Journal*, 63(6), 2301494. <https://doi.org/10.1183/13993003.01494-2023>
- Baldassarri, S. R. (2020). Electronic Cigarettes: Past, Present, and Future: What Clinicians Need to Know. *Clinics in Chest Medicine*, 41(4), 797–807. <https://doi.org/10.1016/j.ccm.2020.08.018>
- Barrera-Núñez, D. A., Zavala-Arciniega, L., Austria-Corrales, F., Romero, M., Barrientos-Gutiérrez, I., Tenorio-Bello, A., & Reynales-Shigematsu, L. M. (2025). Tendencias de uso exclusivo y dual de tabaco combustible y cigarro electrónico en población mexicana. *Salud Pública de México*, 67(3), 276–281. <https://doi.org/10.21149/16293>
- Centros de Integración Juvenil (CIJ). (2025a). *Reporte de Información Epidemiológica del Consumo de Drogas. Consumo de drogas en solicitantes de tratamiento en Centros de Integración Juvenil. Usuarios de drogas ilícitas. Primer semestre del 2025*. CIJ. <http://www.cij.gob.mx/programas/Investigacion/Epidemiologica.asp>
- Centros de Integración Juvenil. (CIJ) (2025b). Sondeo de factores de riesgo del uso de drogas con estudiantes de escuelas de nivel medio y medio superior en México. [Comunicación oficial]
- Czaplicki, L., Kostygina, G., Kim, Y., Perks, S. N., Szczypka, G., Emery, S. L., Vallone, D., & Hair, E. C. (2020). Characterising JUUL-related posts on Instagram. *Tobacco Control*, 29(6), 612–617. <https://doi.org/10.1136/tobaccocontrol-2018-054824>
- Díaz, M. C., Silver, N. A., Bertrand, A., & Schillo, B. A. (2025). Bigger, stronger, and cheaper: growth in e-cigarette market driven by disposable devices with more e-liquid, higher nicotine concentration and declining prices. *Tobacco Control*, 34(1), 65–70. <https://doi.org/10.1136/tc-2023-058033>
- Feeney, S., Rossetti, V., & Terrien, J. (2022). E-Cigarettes—a review of the evidence—harm versus harm reduction. *Tobacco Use Insights*, 15. <https://doi.org/10.1177/1179173X221087524>
- Gaiha, S. M., Lempert, L. K., Lung, H., Vescia, F., & Halpern-Felsher, B. (2024). Appealing characteristics of E-cigarette marketing in the retail environment among adolescents. *Preventive Medicine Reports*, 43, 102769. <https://doi.org/10.1016/j.pmedr.2024.102769>
- Glantz, S. A., Nguyen, N., & Oliveira, A. L. (2024). Population-Based Disease Odds for E-Cigarettes and Dual Use versus Cigarettes. *NEJM Evidence*, 3(3). <https://doi.org/10.1056/EVIDoa2300229>
- Golder, S., Hartwell, G., Barnett, L. M., Nash, S. G., Petticrew, M., & Glover, R. E. (2025). Vaping and harm in young people: umbrella review. *Tobacco Control*. <https://doi.org/10.1136/tc-2024-059219>
- Guraka, A., Mierlea, S., Drake, S. J., Shawa, I. T., Waldron, J., Corcoran, M., Dowse, D., Walkman, D., Burn, L., Sivasubramanian, S., & Kermanizadeh, A. (2024). A comprehensive toxicological analysis of panel of unregulated e-cigarettes to human health. *Toxicology*, 509, 153964. <https://doi.org/10.1016/j.tox.2024.153964>
- Hariri, L. P., Flashner, B. M., Kanarek, D. J., O'Donnell, W. J., Soskis, A., Ziehr, D. R., Frank, A., Nandy, S., Berigei, S. R., Sharma, A., Mathisen, D., Keyes, C. M., Lanuti, M., Muniappan, A., Shepard, J. O., Mino-Kenudson, M., Ly, A., Hung, Y. P., Castellino, F. V., (...) & Christiani, D. C. (2022). E-cigarette use, small airway fibrosis, and constrictive bronchiolitis. *The New England Journal of Medicine Evidence*, 1(6). <https://doi.org/10.1056/EVIDoa2100051>
- Harris, T. (2024). Physical and Chemical Characterization of Aerosols Produced from Experimentally Designed Nicotine Salt-Based E-Liquids. *Chemical Research in Toxicology*, 37(8), 1315–1328. <https://doi.org/10.1021/acs.chemrestox.4c00073>
- Hatsukami, D. K., & Carroll, D. M. (2020). Tobacco harm reduction: Past history, current controversies, and a proposed approach for the future. *Preventive Medicine*, 140, 106099. <https://doi.org/10.1016/j.ypmed.2020.106099>
- Izquierdo-Condoy, J. S., Ruiz, K., Salazar-Santoliva, C., Restrepo, N., Olaya-Villareal, G., Castillo-Concha, J. S., Loaiza-Guevara, V., & Ortiz-Prado, E. (2024). E-cigarette use among adolescents in Latin America: A systematic review of prevalence and associated factors. *Preventive Medicine Reports*, 49, 102952. <https://doi.org/10.1016/j.pmedr.2024.102952>
- Jackler, R. K., Chau, C., Getachew, B. D., Whitcomb, M. M., Lee-Heidenreich, J., Bhatt, A. M., Kim-O'Sullivan, S. H. S., Hoffman, Z. A., Jackler, L. M., & Ramamurthi, D. (2019). *JUUL Advertising Over its First Three Years on the Market*. Stanford University School of Medicine. https://tobacco-img.stanford.edu/wp-content/uploads/2021/07/21231836/JUUL_Marketing_Stanford.pdf
- Javed, S., Usmani, S., Sarfraz, Z., Sarfraz, A., Hanif, A., Firoz, A., Baig, R., Sharath M., Walia, N., Chérrez-Ojeda, I., & Ahmed, S. (2022). A Scoping Review of Vaping, E-Cigarettes, and Mental Health Impact: Depression and Suicidality. *Journal of Community Hospital Internal Medicine Perspectives*, 12(3), 33–39. <https://doi.org/10.55729/2000-9666.1053>
- Levett, J. Y., Filion, K. B., Reynier, P., Prell, C., & Eisenberg, M. J. (2023). Efficacy and Safety of E-Cigarette Use for Smoking Cessation: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *The American Journal of Medicine*, 136(8), 804–813. <https://doi.org/10.1016/j.amjmed.2023.04.014>
- Meehan, J., Heffron, M., McAvoy, H., Reynolds, C., Kyne, L., & Cox, D. W. (2024). The adverse effects of vaping in young people. *Global Pediatrics*, 9, 100190. <https://doi.org/10.1016/j.gped.2024.100190>
- Miller, D. R., Buettner-Schmidt, K., Orr, M., Rykal, K., & Niewojna, E.

- (2021). A systematic review of refillable e-liquid nicotine content accuracy. *Journal of the American Pharmacists Association*, 61(1), 20-26. <https://doi.org/10.1016/j.japh.2020.09.006>
- Novak, M. L., & Wang, G. Y. (2024). The effect of e-cigarettes on cognitive function: a scoping review. *Psychopharmacology*, 241(7), 1287-1297. <https://doi.org/10.1007/s00213-024-06607-8>
- Omaiye, E. E., Cordova, I., Davis, B., & Talbot, P. (2017). Counterfeit Electronic Cigarette Products with Mislabeled Nicotine Concentrations. *Tobacco Regulatory Science*, 3(3), 347-357. <https://doi.org/10.18001/TRS.3.3.10>
- Organización Mundial de la Salud (OMS). (2016). *Sistemas electrónicos de administración de nicotina y sistemas similares sin nicotina (Informe FCTC/COP/7/11)*. OMS. <https://fctc.who.int/docs/librariesprovider12/meeting-reports/fctc-cop-7-11-es.pdf>
- Organización Mundial de la Salud (OMS). (2020). *Electronic nicotine and non-nicotine delivery systems: a brief*. OMS. <https://iris.who.int/server/api/core/bitstreams/ae5bfd99-3a25-4fc3-ac12-7f2e24c60ac5/content>
- Organización Mundial de la Salud (OMS). (2023). *Hay que actuar con urgencia para evitar que los niños y los jóvenes consuman cigarrillos electrónicos* [Comunicado de Prensa]. OMS. <https://www.who.int/es/news/item/14-12-2023-urgent-action-needed-to-protect-children-and-prevent-the-uptake-of-e-cigarettes>
- Organización Mundial de la Salud (OMS). (2024a). *Electronic cigarettes: Call to action*. OMS. <https://cdn.who.int/media/docs/default-source/tobacco-hq/regulating-tobacco-products/ends-call-to-action.pdf>
- Organización Mundial de la Salud (OMS). (2024b). *Enganchando a la próxima generación: cómo capta la industria tabacalera a los clientes jóvenes*. OMS. <https://iris.who.int/handle/10665/378116>
- Organización Mundial de la Salud (OMS). (2025). *WHO global report on trends in prevalence of tobacco use 2000-2024 and projections 2025-2030*. OMS. <https://www.who.int/publications/item/9789240116276>
- Pennington, C., & Hernandez, S. (2025). Disposable electronic cigarettes: Chemical composition in new and used devices. *Journal of Chromatography A*, 1758, 466178. <https://doi.org/10.1016/j.chroma.2025.466178>
- Peri, K., & Eisenberg, M. J. (2024). Feasibility of restricting e-cigarettes to prescription only for smoking cessation. *Respiratory Research*, 25(1). <https://doi.org/10.1186/s12931-024-02792-4>
- Przulj, D., Pesola, F., Myers Smith, K., McRobbie, H., Coleman, T., Lewis, S., Griffith, C., Walton, R., Whitmore, R., Clark, M., Ussher, M., Sinclair, L., Seager, E., Cooper, S., Bauld, L., Naughton, F., Sasieni, P., Manyonda, I., & Hajek, P. (2023). Helping pregnant smokers quit: a multi-centre randomised controlled trial of electronic cigarettes versus nicotine replacement therapy. *Health Technology Assessment* 27(13), 1-53. <https://doi.org/10.3310/AGTH6901>
- Reynolds, L. M., Faure, P., & Barik, J. (2025). Adolescent nicotine exposure and persistent neurocircuitry changes: unveiling lifelong psychiatric risks. *Molecular Psychiatry*, 30. <https://doi.org/10.1038/s41380-025-03110-0>
- Schmidt, C. (2024). Nicotine, Flavor, and More: E-Cigarette Aerosols Deliver Toxic Metals. *Environmental Health Perspectives*, 132(2). <https://doi.org/10.1289/EHP14425>
- Starnini, J. G., Natalello, G., Nigroli, F., Diana, C., Bargagli, E., & Melani, A. S. (2025). Electronic Nicotine Delivery Systems (ENDS): Implications for the Clinician. *Pulmonary Therapy*, 11, 387-404. <https://doi.org/10.1007/s41030-025-00305-2>
- Wakefield, T., Bialous, S., Ling, P., & Apollonio, D. E. (2025). Efforts to Secure Nicotine and Cannabis Product Placements in Popular Media by Ploom, Pax, and Juul: An Analysis of Tobacco Industry Documents. *Nicotine & Tobacco Research*. <https://doi.org/10.1093/ntr/ntaf180>
- Williams, M., & Talbot, P. (2019). Design Features in Multiple Generations of Electronic Cigarette Atomizers. *International Journal of Environmental Research and Public Health*, 16(16), 2904. <https://doi.org/10.3390/ijerph16162904>